

# A Ulm-like Method for Inverse Singular Value Problems

Seak-weng Vong\*

Zheng-jian Bai<sup>†</sup>

Xiao-qing Jin<sup>‡</sup>

## Abstract

In this paper, a Ulm-like method is proposed for solving inverse singular value problems. This method has an advantage over Newton's methods since it avoids solving approximate Jacobian equations. Under some mild assumptions, we show that the proposed method converges at least quadratically in the root sense. Our numerical tests, based on comparison with the inexact Newton method given by Bai and Xu [*Linear Algebra Appl.*, 429 (2008), pp. 527–547], demonstrate the effectiveness of the new method.

**Keywords.** Inverse singular value problem, the Newton method, the Ulm-like method

**AMS subject classification.** 65F18, 65F10, 15A18

## 1 Introduction

Inverse eigenvalue problems (IEPs) arise in various applications such as geophysics, control design, exploration and remote sensing, principal component analysis, molecular spectroscopy, particle physics, circuit theory, and applied mechanics, etc. One may refer to [1, 2, 6, 9, 10, 11, 12, 22] for the applications, mathematical theory, and algorithmic aspects of general IEPs. As a natural extension of IEPs, inverse singular value problems (ISVPs) also have a growing importance in practical applications including the optimal sequence design for direct-spread code division multiple access [20] and the construction of nonnegative and positive matrices from given singular values [14, 15].

In this paper, we consider the following inverse singular value problem.

**ISVP** Given  $n + 1$  real  $m$ -by- $n$  matrices  $A_0, A_1, \dots, A_n$  with  $m \geq n$ , and  $n$  nonnegative real numbers with an order  $\sigma_1^* \geq \sigma_2^* \geq \dots \geq \sigma_n^*$ , find a vector  $\mathbf{c} = (c_1, \dots, c_n)^T \in \mathbb{R}^n$  such that  $\{\sigma_j^*\}_{j=1}^n$  are exactly the singular values of the matrix  $A(\mathbf{c})$  defined by

$$A(\mathbf{c}) \equiv A_0 + c_1 A_1 + c_2 A_2 + \dots + c_n A_n. \quad (1)$$

The ISVP was first addressed by Chu in [8], where the author gave a continuous approach and an iterative approach for solving the ISVP. In particular, the iterative approach is actually

---

\*Department of Mathematics, University of Macau, Macao, P. R. China (swvong@umac.mo).

<sup>†</sup>School of Mathematical Sciences, Xiamen University, Xiamen 361005, P. R. China (zjbai@xmu.edu.cn). The research of this author is partially supported by the Natural Science Foundation of Fujian Province of China for Distinguished Young Scholars (No. 2010J06002).

<sup>‡</sup>Department of Mathematics, University of Macau, Macao, P. R. China (xqjin@umac.mo). The research of this author is supported by the grant UL020/08-Y3/MAT/JXQ01/FST from University of Macau.

Newton’s method which generalizes a numerical method proposed by Friedland, Nocedal, and Overton [12] for solving a kind of inverse eigenvalue problem.

In this paper, we propose a Ulm-like method for solving the ISVP. This is motivated by four papers due to [3, 5, 18, 19]. In [5], an inexact Newton-type method was given for solving the ISVP. In [3, 18, 19], Ulm-like methods (e.g., [21]) were presented for solving IEPs. Ulm’s method is an iterative method like Newton’s method, which is originally proposed for solving a nonlinear equation of  $\mathbf{g}(\mathbf{x}) = \mathbf{0}$ , where  $\mathbf{g}$  is a Fréchet-differentiable operator defined on a convex subset  $\mathcal{D}$  of a Banach space  $\mathcal{X}$  with values in another Banach space  $\mathcal{Y}$ . The Ulm method generates a sequence of

$$\mathbf{x}_{k+1} = \mathbf{x}_k - Q_k \mathbf{g}(\mathbf{x}_k), \quad Q_{k+1} = Q_k + (\mathcal{I} - Q_k \mathbf{g}'(\mathbf{x}_{k+1})) Q_k, \quad k = 0, 1, \dots \quad (2)$$

Here,  $\mathcal{I}$  denotes the identity operator,  $\mathbf{g}'(\mathbf{x}) \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$  the space of bounded linear operators from  $\mathcal{X}$  into  $\mathcal{Y}$ , and  $Q_k \in \mathcal{L}(\mathcal{Y}, \mathcal{X})$  ( $k \geq 0$ ). The Ulm method (2) generates a sequence  $\{\mathbf{x}_k\}$  which converges to a locally unique solution  $\mathbf{x}^*$  of  $\mathbf{g}(\mathbf{x}) = \mathbf{0}$  at least quadratically in the root sense. For the definition of convergence rate in the root sense, see [16, Chapter 9] or Definition 3.8 in Section 3. Moreover, the method provides successive approximation  $Q_k$  to  $\mathbf{g}'(\mathbf{x}^*)^{-1}$  simultaneously.

Our proposed Ulm-like method can avoid solving approximate Jacobian equations appeared in the inexact Newton-type method and thus reduce the complexity caused by possible ill-conditioned Jacobian equations. Under some mild assumptions, we show that our method converges at least quadratically in the root sense. Numerical tests demonstrate that the proposed method is very effective.

Throughout the paper, we use the following notations. Let  $\|\cdot\|$  be the Euclidean vector norm or its induced matrix norm and  $\|\cdot\|_F$  denote the Frobenius norm. We use  $I$  to denote an identity matrix of appropriate size. For any  $\mathbf{c} \in \mathbb{R}^n$ ,  $\{\sigma_j(A(\mathbf{c}))\}_{j=1}^n$ ,  $\{\mathbf{u}_j(A(\mathbf{c}))\}_{j=1}^m$ , and  $\{\mathbf{v}_j(A(\mathbf{c}))\}_{j=1}^n$  stand for the singular values, the left singular vectors, and the right singular vectors of  $A(\mathbf{c})$ , respectively. Let  $\boldsymbol{\sigma}^* = (\sigma_1^*, \dots, \sigma_n^*)^T \in \mathbb{R}^n$  and  $\Sigma^* = \text{diag}(\sigma_1^*, \dots, \sigma_n^*) \in \mathbb{R}^{m \times n}$ .

The paper is organized as follows. In Section 2, we present the Ulm-like method for the ISVP. In Section 3, we show our Ulm-like method converges at least quadratically in the root sense. Numerical tests are reported in Section 4 and some concluding remarks are given in Section 5.

## 2 A Ulm-like Method

In this section, we propose a Ulm-like method for solving the ISVP. We note that solving the ISVP is equivalent to finding a solution to the nonlinear equation

$$\mathbf{f}(\mathbf{c}) \equiv \boldsymbol{\sigma}(A(\mathbf{c})) - \boldsymbol{\sigma}^* = \mathbf{0}, \quad \boldsymbol{\sigma}(A(\mathbf{c})) = (\sigma_1(A(\mathbf{c})), \dots, \sigma_n(A(\mathbf{c})))^T. \quad (3)$$

Thus we can use the Ulm method (2) to solve (3). In contrast to the Newton method [8] and the inexact Newton method [5], where approximate Jacobian equations need to be solved, the Ulm method (2) is in “inversion free” and keeps away from the solution of Jacobian equations. Moreover, successive approximations  $Q_k$  to  $\mathbf{g}'(\mathbf{x}^*)$  may be useful for the subsequent convergence analysis. The Ulm method (2) was successfully employed in solving IEPs [3, 18, 19]. This motivates us to propose the following Ulm-like method for the ISVP.

**Algorithm 1: The Ulm-like Method**

- I. Given  $\mathbf{c}^0 \in \mathbb{R}^n$ , compute the normalized left singular vectors  $\{\mathbf{u}_i^0 = \mathbf{u}_i(A(\mathbf{c}^0))\}_{i=1}^m$ , and the normalized right singular vectors  $\{\mathbf{v}_i^0 = \mathbf{v}_i(A(\mathbf{c}^0))\}_{i=1}^n$  of  $A(\mathbf{c}^0)$ . Form Jacobian matrix  $J_0$  and  $\mathbf{w}^0 \in \mathbb{R}^n$  by

$$[J_0]_{ij} = (\mathbf{u}_i^0)^T A_j \mathbf{v}_i^0, \quad [\mathbf{w}^0]_i = (\mathbf{u}_i^0)^T A_0 \mathbf{v}_i^0, \quad i, j = 1, \dots, n. \quad (4)$$

Let  $Q_0 = J_0^{-1}$  and compute  $\mathbf{c}^1$  by

$$\mathbf{c}^1 = \mathbf{c}^0 - Q_0(J_0 \mathbf{c}^0 + \mathbf{w}^0 - \boldsymbol{\sigma}^*) = J_0^{-1}(\boldsymbol{\sigma}^* - \mathbf{w}^0).$$

Set  $\mathbf{s}^0 \equiv (s_1^0, s_2^0, \dots, s_n^0)^T = \boldsymbol{\sigma}^*$ .

- II. For  $k = 1, 2, \dots$ , until convergence, do:

(i) Form the matrix  $A(\mathbf{c}^k)$  by (1).

(ii) Form the matrix  $Z_{k-1} \equiv U_{k-1}^T A(\mathbf{c}^k) V_{k-1}$ .

(iii) Calculate the skew-symmetric matrices  $X_k$  and  $Y_k$  by

$$\begin{aligned} [X_k]_{ij} &= 0, \quad n+1 \leq i \neq j \leq m; \\ [X_k]_{ij} &= -[X_k]_{ji} = \frac{[Z_{k-1}]_{ij}}{s_j^{k-1}}, \quad n+1 \leq i \leq m, \quad 1 \leq j \leq n; \\ [X_k]_{ij} &= -[X_k]_{ji} = \frac{s_i^{k-1}[Z_{k-1}]_{ji} + s_j^{k-1}[Z_{k-1}]_{ij}}{(s_j^{k-1})^2 - (s_i^{k-1})^2}, \quad 1 \leq i < j \leq n; \\ [Y_k]_{ij} &= -[Y_k]_{ji} = \frac{s_i^{k-1}[Z_{k-1}]_{ij} + s_j^{k-1}[Z_{k-1}]_{ji}}{(s_j^{k-1})^2 - (s_i^{k-1})^2}, \quad 1 \leq i < j \leq n. \end{aligned}$$

(iv) Compute  $U_k = [\mathbf{u}_1^k, \dots, \mathbf{u}_m^k]$  and  $V_k = [\mathbf{v}_1^k, \dots, \mathbf{v}_n^k]$  by solving

$$\left(I + \frac{1}{2}X_k\right)U_k^T = \left(I - \frac{1}{2}X_k\right)U_{k-1}^T, \quad \left(I + \frac{1}{2}Y_k\right)V_k^T = \left(I - \frac{1}{2}Y_k\right)V_{k-1}^T.$$

(v) Form the approximate Jacobian matrix  $J_k$  and  $\mathbf{w}^k \in \mathbb{R}^n$  by

$$[J_k]_{ij} = (\mathbf{u}_i^k)^T A_j \mathbf{v}_i^k, \quad [\mathbf{w}^k]_i = (\mathbf{u}_i^k)^T A_0 \mathbf{v}_i^k, \quad i, j = 1, \dots, n.$$

(vi) Compute  $Q_k \in \mathbb{R}^{n \times n}$  and  $\mathbf{c}^{k+1}$  by

$$Q_k = Q_{k-1} + (I - Q_{k-1}J_k)Q_{k-1}, \quad \mathbf{c}^{k+1} = \mathbf{c}^k - Q_k(J_k \mathbf{c}^k + \mathbf{w}^k - \boldsymbol{\sigma}^*).$$

Set  $\mathbf{s}^k \equiv (s_1^k, s_2^k, \dots, s_n^k)^T = \boldsymbol{\sigma}^* + \mathbf{t}^k$ , where

$$\mathbf{t}^k \equiv (t_1^k, t_2^k, \dots, t_n^k)^T = (I - J_k Q_k)(J_k \mathbf{c}^k + \mathbf{w}^k - \boldsymbol{\sigma}^*).$$

We note that, instead of solving successive approximate Jacobian equations, Algorithm 1 generates successive approximations  $Q_k$  to the inverse of Jacobian matrix  $J(\mathbf{c}^*)$  (see (8) below for the definition), provided that a solution  $\mathbf{c}^*$  to the ISVP exists. We observe from Algorithm 1 that two orthogonal matrices  $U_k$  and  $V_k$  are given by

$$U_k = U_{k-1}\Phi_k, \quad V_k = V_{k-1}\Psi_k, \quad (5)$$

where  $\Phi_k = (I + \frac{1}{2}X_k)(I - \frac{1}{2}X_k)^{-1}$  and  $\Psi_k = (I + \frac{1}{2}Y_k)(I - \frac{1}{2}Y_k)^{-1}$ . For Algorithm 1, it is easy to check that the vector  $\mathbf{c}^{k+1}$  and the skew-symmetric matrices  $X_{k+1}$  and  $Y_{k+1}$  are determined by

$$S_k + X_{k+1}S_k - S_kY_{k+1} = U_k^T A(\mathbf{c}^{k+1})V_k, \quad (6)$$

where  $S_k = \Sigma^* + T_k$  with  $S_k = \text{diag}(s_1^k, \dots, s_n^k) \in \mathbb{R}^{m \times n}$  and  $T_k = \text{diag}(t_1^k, \dots, t_n^k) \in \mathbb{R}^{m \times n}$ . We should mention that the operation cost of Algorithm 1 is  $O(n^3)$ , almost the same as that of solving Jacobian equation. However, computing the product of matrices is simpler than solving equations and has no un-stability problem caused by ill-conditioning in solving equations. In particular, the parallel computation techniques can be applied in the product of matrices in the Ulm-like method to improve the computational efficiency. We will show that Algorithm 1 converges at least quadratically in the root sense in next section.

### 3 Convergence Analysis

In this section, we establish a quadratic convergence in the root sense of Algorithm 1 for the ISVP. Suppose that the given singular values  $\{\sigma_j^*\}_{j=1}^n$  are all positive and distinct. As noted in [7], the ISVP may not have a solution since singular values cannot be assigned arbitrarily. Therefore, in what follows, we assume that the ISVP has a solution  $\mathbf{c}^*$ . Let  $A(\mathbf{c}^*) = U^*\Sigma^*(V^*)^T$  be the singular value decomposition of  $A(\mathbf{c}^*)$ , where  $U^* \in \mathcal{O}(m)$  and  $V^* \in \mathcal{O}(n)$ . It follows from Theorem 1.9.3 in [22] that there exists a neighborhood  $\mathcal{B}(\mathbf{c}^*)$  at  $\mathbf{c}^*$  such that the singular values  $\sigma_j(A(\mathbf{c}))$  are all distinct and differentiable for all  $\mathbf{c} \in \mathcal{B}(\mathbf{c}^*)$ . For any  $\mathbf{c} \in \mathcal{B}(\mathbf{c}^*)$ , the function  $\mathbf{f}(\mathbf{c})$  defined by (3) is nonlinear and continuously differentiable. For any  $\mathbf{c} \in \mathcal{B}(\mathbf{c}^*)$ , by using

$$\sigma_i(A(\mathbf{c})) = \mathbf{u}_i(A(\mathbf{c}))^T A(\mathbf{c}) \mathbf{v}_i(A(\mathbf{c})), \quad \mathbf{u}_i(A(\mathbf{c}))^T \mathbf{u}_i(A(\mathbf{c})) = 1, \quad \mathbf{v}_i(A(\mathbf{c}))^T \mathbf{v}_i(A(\mathbf{c})) = 1, \quad (7)$$

it is easy to derive that

$$\frac{\partial \sigma_i(A(\mathbf{c}))}{\partial c_j} = \mathbf{u}_i(A(\mathbf{c}))^T A_j \mathbf{v}_i(A(\mathbf{c})).$$

Then we get Jacobian matrix  $J(\mathbf{c})$  of  $\mathbf{f}$  at  $\mathbf{c} \in \mathcal{B}(\mathbf{c}^*)$ , where

$$[J(\mathbf{c})]_{ij} = \mathbf{u}_i(A(\mathbf{c}))^T A_j \mathbf{v}_i(A(\mathbf{c})), \quad 1 \leq i, j \leq n. \quad (8)$$

We obtain by (7) and (8),

$$\boldsymbol{\sigma}(A(\mathbf{c})) = J(\mathbf{c})\mathbf{c} + \mathbf{w}(\mathbf{c}), \quad (9)$$

where

$$\boldsymbol{\sigma}(A(\mathbf{c})) = (\sigma_1(A(\mathbf{c})), \dots, \sigma_n(A(\mathbf{c})))^T, \quad [\mathbf{w}(\mathbf{c})]_i = \mathbf{u}_i(A(\mathbf{c}))^T A_0 \mathbf{v}_i(A(\mathbf{c})), \quad 1 \leq i \leq n.$$

To show the convergence of Algorithm 1, in what follows, we assume that Jacobian matrix  $J(\mathbf{c}^*)$  defined by (8) is nonsingular. The continuity of the matrix and its inverse ensures there exist a scalar  $\delta_1 > 0$  and a constant  $C$  such that if

$$\max\{\|[\mathbf{u}_1, \dots, \mathbf{u}_n] - U_1^*\|, \|[\mathbf{v}_1, \dots, \mathbf{v}_n] - V^*\|\} \leq \delta_1,$$

then the approximate Jacobian matrix  $J = [\mathbf{u}_i^T A_j \mathbf{v}_i]$  is nonsingular and  $\|J^{-1}\| \leq C$ . Also, let the matrices  $\Sigma^*$ ,  $U^*$ , and  $U_k$  be partitioned as follows:

$$\Sigma^* = \begin{bmatrix} \Sigma_1^* \\ 0 \end{bmatrix}, \quad \Sigma_1^* \in \mathbb{R}^{n \times n}; \quad U^* = [U_1^*, U_2^*], \quad U_1^* \in \mathbb{R}^{m \times n}; \quad U_k = [U_1^k, U_2^k], \quad U_1^k \in \mathbb{R}^{m \times n}.$$

### 3.1 Preliminary Results

We recall some necessary preliminary lemmas which can be found in [4, 5, 13].

**Lemma 3.1** [13, Corollary 8.6.2] *Let  $M, M + \Delta M \in \mathbb{R}^{m \times n}$  with  $m \geq n$ . For any  $1 \leq k \leq n$ ,*

$$|\sigma_k(M + \Delta M) - \sigma_k(M)| \leq \|\Delta M\|,$$

where  $\sigma_k(M)$  is the  $k$ th largest singular value of  $M$ .

**Lemma 3.2** [4, Lemma 2] *For any  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ , we have*

$$\|A(\mathbf{x}) - A(\mathbf{y})\| \leq \zeta \|\mathbf{x} - \mathbf{y}\|,$$

where  $\zeta = \left( \sum_{i=1}^n \|A_i\|^2 \right)^{1/2}$ .

**Lemma 3.3** [4, Lemma 4] *Suppose that the given singular values  $\{\sigma_j^*\}_{j=1}^n$  are all positive and distinct. Then there exist positive numbers  $\delta_2$  and  $\kappa$  such that, when  $\|\mathbf{c}^k - \mathbf{c}^*\| \leq \delta_2$ ,*

$$\begin{aligned} \|[\mathbf{u}_1(A(\mathbf{c}^k)), \dots, \mathbf{u}_n(A(\mathbf{c}^k))] - U_1^*\| &\leq \kappa \|\mathbf{c}^k - \mathbf{c}^*\|, \\ \|[\mathbf{v}_1(A(\mathbf{c}^k)), \dots, \mathbf{v}_n(A(\mathbf{c}^k))] - V^*\| &\leq \kappa \|\mathbf{c}^k - \mathbf{c}^*\|. \end{aligned}$$

**Lemma 3.4** [4, Lemma 5] *Let  $B \in \mathbb{R}^{n \times n}$  such that  $\|B\| < 1$ . Then  $I - \frac{1}{2}B$  is nonsingular and*

$$\left\| \left( I + \frac{1}{2}B \right) \left( I - \frac{1}{2}B \right)^{-1} - (I + B) \right\| \leq \|B\|^2.$$

We can easily obtain the following result based on Lemma 6 in [4].

**Lemma 3.5** *Let  $Z \in \mathbb{R}^{m \times n}$  and  $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_n) \in \mathbb{R}^{m \times n}$  with  $m \geq n$ , where  $\sigma_1 > \sigma_2 > \dots > \sigma_n > 0$ . Suppose that two skew-symmetric matrices  $X \in \mathbb{R}^{m \times n}$  and  $Y \in \mathbb{R}^{n \times n}$  satisfy*

$$X\Sigma - \Sigma Y = Z.$$

Then we have

$$\|X\| \leq \left( \frac{2n\sigma_1}{d} + \frac{1}{\sigma_n} \right) \|Z\|, \quad \|Y\| \leq \frac{2n\sigma_1}{d} \|Z\|,$$

where  $d = \min_{i \neq j} |\sigma_i^2 - \sigma_j^2|$ .

### 3.2 Convergence Results

Let

$$\begin{aligned}
\sigma_0^* &= 2\sigma_1^*, & \sigma_{n+1}^* &= \frac{1}{2}\sigma_n^*, \\
d^* &= \min_{1 \leq i \neq j \leq n} |(\sigma_i^*)^2 - (\sigma_j^*)^2|, & \widehat{d}^* &= \min_{1 \leq j \leq n+1} |\sigma_{j-1}^* - \sigma_j^*|, \\
\rho &= \frac{6n\sigma_1^*}{d^*} + \frac{4}{3\sigma_n^*}, & \mu_0 &= 2(8nC \max_j \|A_j\|)^2, \\
\mu_1 &= \sqrt{2} \rho (\zeta^2 C \sqrt{n} + \zeta), & \mu_2 &= 23\sigma_1^*, \\
\mu_3 &= (2C + 1)\sqrt{n}(\mu_2 + \sqrt{n}), & \mu_4 &= \sqrt{2}\rho[(\zeta + 1)\mu_3 + 2\mu_2 + \sqrt{n}], \\
\mu &= \max\{\mu_0 + 2, \mu_2 + \sqrt{n}, \mu_3, \mu_4\}.
\end{aligned}$$

We give the following result for the initial step of Algorithm 1.

**Theorem 3.6** *There exists a scalar  $\delta_3 > 0$  such that, when  $\|\mathbf{c}^0 - \mathbf{c}^*\| \leq \delta_3$ ,*

$$\max\{\|U_1^0 - U_1^*\|, \|V_0 - V^*\|\} \leq \kappa \|\mathbf{c}^0 - \mathbf{c}^*\| \leq \delta_1/4,$$

$$\sqrt{\|X_1\|^2 + \|Y_1\|^2} \leq \mu_1 \|\mathbf{c}^0 - \mathbf{c}^*\| \leq \delta_1/4.$$

**Proof:** Let  $\|\mathbf{c}^0 - \mathbf{c}^*\| \leq \delta_3$ , where

$$\delta_3 = \min \left\{ 1, \delta_2, \frac{\delta_1}{4\kappa}, \frac{\delta_1}{4\mu_1} \right\}.$$

By Lemma 3.3, we have

$$\max\{\|U_1^0 - U_1^*\|, \|V_0 - V^*\|\} \leq \kappa \|\mathbf{c}^0 - \mathbf{c}^*\| \leq \delta_1/4.$$

Thus Jacobian matrix  $J_0$  is invertible and  $\|J_0^{-1}\| \leq C$ . By Lemmas 3.1 and 3.2, we obtain

$$\max_j |\sigma_j(A(\mathbf{c}^0)) - \sigma_j^*| \leq \|A(\mathbf{c}^0) - A(\mathbf{c}^*)\| \leq \zeta \|\mathbf{c}^0 - \mathbf{c}^*\|. \quad (10)$$

From Algorithm 1, (9), and (10), it follows that

$$\begin{aligned}
\|\mathbf{c}^1 - \mathbf{c}^0\| &\leq \|Q_0\| \cdot \|J_0 \mathbf{c}^0 + \mathbf{w}^0 - \boldsymbol{\sigma}^*\| = \|Q_0\| \cdot \|J(\mathbf{c}^0) \mathbf{c}^0 + \mathbf{w}(\mathbf{c}^0) - \boldsymbol{\sigma}^*\| \\
&\leq C \|\boldsymbol{\sigma}(A(\mathbf{c}^0)) - \boldsymbol{\sigma}^*\| \leq C \sqrt{n} \max_j |\sigma_j(A(\mathbf{c}^0)) - \sigma_j^*| \leq \zeta C \sqrt{n} \|\mathbf{c}^0 - \mathbf{c}^*\|.
\end{aligned} \quad (11)$$

Let  $\Sigma_0 = \text{diag}[\sigma_1(A(\mathbf{c}^0)), \dots, \sigma_n(A(\mathbf{c}^0))] \in \mathbb{R}^{m \times n}$ . Then

$$U_0^T A(\mathbf{c}^0) V_0 = \Sigma_0. \quad (12)$$

We have by (6) and  $S_0 = \Sigma^*$ ,

$$U_0^T A(\mathbf{c}^1) V_0 = S_0 + X_1 S_0 - S_0 Y_1 = \Sigma^* + X_1 \Sigma^* - \Sigma^* Y_1. \quad (13)$$

By using (12) and (13), we obtain

$$X_1 \Sigma^* - \Sigma^* Y_1 = U_0^T (A(\mathbf{c}^1) - A(\mathbf{c}^0)) V_0 - (\Sigma^* - \Sigma_0). \quad (14)$$

By Lemmas 3.5 and 3.2, it follows from (10), (11), and (14) that

$$\begin{aligned} \sqrt{\|X_1\|^2 + \|Y_1\|^2} &\leq \sqrt{2}\rho(\|A(\mathbf{c}^1) - A(\mathbf{c}^0)\| + \|\Sigma^* - \Sigma_0\|) \\ &\leq \sqrt{2}\rho(\zeta\|\mathbf{c}^1 - \mathbf{c}^0\| + \max_j |\sigma_j(A(\mathbf{c}^0)) - \sigma_j^*|) \\ &\leq \sqrt{2}\rho(\zeta^2 C \sqrt{n} + \zeta)\|\mathbf{c}^0 - \mathbf{c}^*\| \\ &= \mu_1 \|\mathbf{c}^0 - \mathbf{c}^*\| \leq \frac{\delta_1}{4}. \end{aligned}$$

□

Now, we offer the essential estimates of the quantities generated by Algorithm 1:

$$\|U_k^T A(\mathbf{c}^k) V_k - \Sigma^*\|, \quad \|\mathbf{c}^{k+1} - \mathbf{c}^k\|, \quad \|X_k\|, \quad \|Y_k\|, \quad \|U_{k+1} - U_k\|, \quad \|V_{k+1} - V_k\|.$$

**Theorem 3.7** *Suppose that  $\sigma_1^* > \sigma_2^* > \dots > \sigma_n^* > 0$  and Jacobian matrix  $J(\mathbf{c}^*)$  is invertible. Let  $\{\mathbf{c}^k\}$ ,  $\{Q_k\}$ ,  $\{U_k\}$ , and  $\{V_k\}$  be the sequences generated by Algorithm 1. Assume  $\psi_k \equiv \sqrt{\|X_k\|^2 + \|Y_k\|^2}$ . Then, there exists a constant  $\delta > 0$  such that, when  $\|\mathbf{c}^0 - \mathbf{c}^*\| < \delta$ , the following inequalities hold for all  $k \geq 1$ :*

$$\|U_k^T A(\mathbf{c}^k) V_k - \Sigma^*\| \leq \mu^{2^k-1} \psi_1^{2^k}, \quad (15)$$

$$\|I - J_k Q_k\| \leq \mu^{2^k-1} \psi_1^{2^k}, \quad (16)$$

$$\|Q_k\| \leq 2C, \quad (17)$$

$$\|\mathbf{c}^{k+1} - \mathbf{c}^k\| \leq \mu^{2^k-1} \psi_1^{2^k}, \quad (18)$$

$$\psi_{k+1} \leq \mu^{2^k-1} \psi_1^{2^k}, \quad (19)$$

$$\|U_{k+1} - U_k\| \leq 2\mu^{2^k-1} \psi_1^{2^k}, \quad (20)$$

$$\|V_{k+1} - V_k\| \leq 2\mu^{2^k-1} \psi_1^{2^k}. \quad (21)$$

Moreover, we have

$$\frac{3}{2}\sigma_1^* > s_1^k > \frac{1}{2}(\sigma_1^* + \sigma_2^*) > s_2^k > \frac{1}{2}(\sigma_2^* + \sigma_3^*) > \dots > \frac{1}{2}(\sigma_{n-1}^* + \sigma_n^*) > s_n^k > \frac{3}{4}\sigma_n^* > 0$$

and

$$\min_{i \neq j} |(s_i^k)^2 - (s_j^k)^2| \geq \frac{1}{2}d^* > 0,$$

which make sure that Algorithm 1 will not break down.

**Proof:** Let  $\|\mathbf{c}^0 - \mathbf{c}^*\| < \delta$  where

$$\delta = \min \left\{ 1, \delta_3, \frac{1}{\mu_1}, \frac{1}{2\mu_1\mu}, \frac{d^*}{2\mu_1\mu(1+4\sigma_1^*)}, \frac{\widehat{d}^*}{2\mu_1\mu}, \frac{\sigma_n^*}{2\mu_1\mu}, \frac{\mu\sigma_n^*}{2\mu_1\sqrt{n}}, \frac{3\delta_1}{16\mu\mu_1} \right\}. \quad (22)$$

We establish our theorem by the induction. We first consider the case of  $k = 1$ .

By (22), we have  $\delta \leq \delta_3$  and  $\delta \leq 1/\mu_1$ . Then by Theorem 3.6,

$$\psi_1 = \sqrt{\|X_1\|^2 + \|Y_1\|^2} < \mu_1\delta \leq \min\{1, \delta_1/4\}, \quad \|U_1^0 - U_1^*\| \leq \delta_1/4, \quad \|V_0 - V^*\| \leq \delta_1/4. \quad (23)$$

Thus by Lemma 3.4 and (23),

$$\|U_1 - U_0\| \leq 2\|X_1\| \leq 2\psi_1 < \frac{\delta_1}{2}, \quad \|V_1 - V_0\| \leq 2\|Y_1\| \leq 2\psi_1 < \frac{\delta_1}{2}. \quad (24)$$

By using (23) and (24), we get

$$\|U_1^1 - U_1^*\| \leq \|U_1^1 - U_1^0\| + \|U_1^0 - U_1^*\| < \frac{\delta_1}{2} + \frac{\delta_1}{4} < \delta_1,$$

and

$$\|V_1 - V^*\| \leq \|V_1 - V_0\| + \|V_0 - V^*\| < \frac{\delta_1}{2} + \frac{\delta_1}{4} < \delta_1.$$

Hence,  $J_1$  is invertible and  $\|J_1^{-1}\| \leq C$ . Furthermore, we have by (5) and Lemma 3.4,

$$\Phi_1 = I + X_1 + G_1, \quad \Psi_1 = I + Y_1 + H_1, \quad (25)$$

where  $\|G_1\| \leq \|X_1\|^2 \leq \psi_1^2$  and  $\|H_1\| \leq \|Y_1\|^2 \leq \psi_1^2$ . Using  $U_1 = U_0\Phi_1$ ,  $V_1 = V_0\Psi_1$ , and (25), one can derive via simple computation,

$$U_1^T A(\mathbf{c}^1) V_1 = S_0 + R_1 = \Sigma^* + R_1, \quad (26)$$

where

$$\begin{aligned} R_1 &= X_1(S_0 - X_1 S_0 + S_0 Y_1) Y_1 - X_1^2 S_0 - S_0 Y_1^2 + G_1^T (S_0 + X_1 S_0 - S_0 Y_1) (I + Y_1) \\ &\quad + (I - X_1 + G_1^T) (S_0 + X_1 S_0 - S_0 Y_1) H_1. \end{aligned}$$

Using (23), (25), and  $S_0 = \Sigma^*$ , we have

$$\begin{aligned} \|R_1\| &\leq 3\sigma_1^* \|X_1\| \cdot \|Y_1\| + \sigma_1^* \|X_1\|^2 + \sigma_1^* \|Y_1\|^2 + 6\sigma_1^* \|X_1\|^2 + 9\sigma_1^* \|Y_1\|^2 \\ &\leq \frac{3}{2}\sigma_1^* (\|X_1\|^2 + \|Y_1\|^2) + 10\sigma_1^* (\|X_1\|^2 + \|Y_1\|^2) \\ &= \frac{23}{2}\sigma_1^* \psi_1^2 \leq \mu_2 \psi_1^2 \leq \mu \psi_1^2, \end{aligned} \quad (27)$$

which verifies that (15) holds for  $k = 1$ .

Notice that

$$I - J_1 Q_1 = I - 2J_1 Q_0 + J_1 Q_0 J_1 Q_0 = (I - J_1 Q_0)^2.$$



Thus

$$\|I - J_1 Q_1\| \leq (\|I - J_0 Q_0\| + \|J_1 - J_0\| \cdot \|Q_0\|)^2 \leq 2\|I - J_0 Q_0\|^2 + 2\|J_1 - J_0\|^2 \|Q_0\|^2. \quad (28)$$

In addition, for  $1 \leq i, j \leq n$ ,

$$\begin{aligned} |[J_1]_{ij} - [J_0]_{ij}| &= |(\mathbf{u}_i^1)^T A_j \mathbf{v}_i^1 - (\mathbf{u}_i^0)^T A_j \mathbf{v}_i^0| \\ &= |(\mathbf{u}_i^1 - \mathbf{u}_i^0)^T A_j \mathbf{v}_i^1 - (\mathbf{u}_i^0)^T A_j (\mathbf{v}_i^0 - \mathbf{v}_i^1)| \\ &\leq \|A_j\| (\|\mathbf{u}_i^1 - \mathbf{u}_i^0\| + \|\mathbf{v}_i^0 - \mathbf{v}_i^1\|) \\ &\leq \|A_j\| (\|U_1 - U_0\| + \|V_1 - V_0\|). \end{aligned} \quad (29)$$

By (24) and (29), we obtain

$$\|J_1 - J_0\| \leq \|J_1 - J_0\|_F \leq n(\|U_1 - U_0\| + \|V_1 - V_0\|) \cdot \max_j \|A_j\| \leq 4n \max_j \|A_j\| \psi_1. \quad (30)$$

By (22), we have  $\delta \leq 1/(2\mu_1\mu)$ . Then, by using (23), (28), (30), and  $Q_0 = J_0^{-1}$ , we get

$$\|I - J_1 Q_1\| \leq 2C^2 (4n \max_j \|A_j\|)^2 \psi_1^2 \leq \mu_0 \psi_1^2 \leq \mu \psi_1^2 < \mu \mu_1 \delta < 1, \quad (31)$$

i.e., (16) holds for  $k = 1$ .

The diagonal entries of (26) yield

$$J_1 \mathbf{c}^1 + \mathbf{w}^1 - \boldsymbol{\sigma}^* = \mathbf{r}_1$$

with  $\mathbf{r}_1$  being the diagonal vector of the matrix  $R_1$ . We therefore have

$$\|\mathbf{c}^2 - \mathbf{c}^1\| \leq \|Q_1\| \cdot \|J_1 \mathbf{c}^1 + \mathbf{w}^1 - \boldsymbol{\sigma}^*\| = \|Q_1\| \cdot \|\mathbf{r}_1\| \leq \sqrt{n} \|Q_1\| \cdot \|R_1\|. \quad (32)$$

Also by (31),

$$\|Q_1\| \leq \|J_1^{-1}\| \cdot \|J_1 Q_1\| \leq \|J_1^{-1}\| (1 + \|I - J_1 Q_1\|) \leq 2C,$$

which shows that (17) holds for  $k = 1$ . Then, we immediately get by (27) and (32),

$$\|\mathbf{c}^2 - \mathbf{c}^1\| \leq 2C \sqrt{n} \mu_2 \psi_1^2 \leq \mu_3 \psi_1^2 \leq \mu \psi_1^2. \quad (33)$$

This confirms that (18) holds for  $k = 1$ .

Recall from (22) that  $\delta \leq 1/(2\mu\mu_1)$  and  $\delta \leq \hat{d}^*/(2\mu\mu_1)$ . We then have by (23), (26), (27), and (31),

$$\begin{aligned} \|\mathbf{s}^1 - \boldsymbol{\sigma}^*\| &\leq \|I - J_1 Q_1\| \cdot \|J_1 \mathbf{c}^1 + \mathbf{w}^1 - \boldsymbol{\sigma}^*\| \\ &\leq \mu_0 \psi_1^2 \sqrt{n} \|R_1\| \leq \sqrt{n} \mu_0 \psi_1^2 \mu_2 \psi_1^2 \\ &\leq \sqrt{n} \mu \psi_1^2 \mu_2 \psi_1^2 \leq \frac{\sqrt{n}}{\mu} (\mu \psi_1)^2 \mu_2 \psi_1^2 \\ &\leq \frac{\sqrt{n}}{\mu} (\mu \mu_1 \delta)^2 \mu_2 \psi_1^2 \leq \mu_2 \psi_1^2 \leq \mu \psi_1^2 \\ &< \mu \mu_1 \delta \leq \min\{\frac{1}{2}, \frac{1}{2} \hat{d}^*\}. \end{aligned} \quad (34)$$

Thus one has

$$\frac{3}{2}\sigma_1^* > s_1^1 > \frac{1}{2}(\sigma_1^* + \sigma_2^*) > s_2^1 > \frac{1}{2}(\sigma_2^* + \sigma_3^*) > \cdots > \frac{1}{2}(\sigma_{n-1}^* + \sigma_n^*) > s_n^1 > \frac{3}{4}\sigma_n^* > 0.$$

By (22), we have  $\delta \leq d^*/(2\mu\mu_1(4\sigma_1^* + 1))$ . Then we further derive from (34) that

$$\begin{aligned} |(s_i^1)^2 - (s_j^1)^2| &\geq d^* - 4\sigma_1^*\|\mathbf{s}^1 - \boldsymbol{\sigma}^*\| - 2\|\mathbf{s}^1 - \boldsymbol{\sigma}^*\|^2 \\ &\geq d^* - (4\sigma_1^* + 1)\|\mathbf{s}^1 - \boldsymbol{\sigma}^*\| \\ &\geq d^* - (4\sigma_1^* + 1)\mu\mu_1\delta \geq \frac{1}{2}d^*, \end{aligned}$$

for  $1 \leq i \neq j \leq n$ .

Notice that  $U_1^T A(\mathbf{c}^2)V_1 = S_1 + X_2S_1 - S_1Y_2$ . Combining this with (26), we obtain

$$X_2S_1 - S_1Y_2 = U_1^T (A(\mathbf{c}^2) - A(\mathbf{c}^1))V_1 + R_1 + (\Sigma^* - S_1).$$

By Lemmas 3.5 and 3.2, it follows from (27), (33), and (34) that

$$\begin{aligned} \psi_2 = \sqrt{\|X_2\|^2 + \|Y_2\|^2} &\leq \sqrt{2}\rho(\|A(\mathbf{c}^2) - A(\mathbf{c}^1)\| + \|R_1\| + \|\Sigma^* - S_1\|) \\ &\leq \sqrt{2}\rho(\zeta\|\mathbf{c}^2 - \mathbf{c}^1\| + \|R_1\| + \|\mathbf{s}^1 - \boldsymbol{\sigma}^*\|) \\ &\leq \sqrt{2}\rho(\zeta\mu_3 + 2\mu_2)\psi_1^2 \leq \mu_4\psi_1^2 \leq \mu\psi_1^2. \end{aligned} \quad (35)$$

This shows (19) holds for  $k = 1$ . In addition, by (22), we have  $\delta \leq 1/(2\mu_1\mu)$ . Then by (23) and (35),

$$\psi_2 \leq \mu\psi_1^2 \leq \mu\mu_1\delta\psi_1 \leq \psi_1 < 1.$$

By Lemma 3.4, it follows from (35) that

$$\|U_2 - U_1\| \leq 2\|X_2\| \leq 2\mu_4\psi_1^2 \leq 2\mu\psi_1^2, \quad \|V_2 - V_1\| \leq 2\|Y_2\| \leq 2\mu_4\psi_1^2 \leq 2\mu\psi_1^2,$$

which shows that (20) and (21) hold for  $k = 1$ .

Now, we consider the general case. Suppose that (15)–(21) hold for all positive integers less than or equal to  $k - 1$ . By (22), we have  $\delta \leq 1/(2\mu_1\mu)$ . Then, by the hypothesis, based on (23), it is easy to check that

$$\psi_k \leq \mu^{2^{k-1}-1}\psi_1^{2^{k-1}} = (\mu\psi_1)^{2^{k-1}-1}\psi_1 \leq (\mu\mu_1\delta)^{2^{k-1}-1}\psi_1 \leq \psi_1 < 1. \quad (36)$$

By the hypothesis again, one has

$$\|I - J_{k-1}Q_{k-1}\| \leq \mu^{2^{k-1}-1}\psi_1^{2^{k-1}} \leq \psi_1 < 1 \quad (37)$$

and

$$\|U_{k-1}^T A(\mathbf{c}^{k-1})V_{k-1} - \Sigma^*\| \leq \mu^{2^{k-1}-1}\psi_1^{2^{k-1}}. \quad (38)$$

By (22) again, we have  $\delta \leq 1/(2\mu\mu_1)$  and  $\delta \leq \widehat{d}^*/(2\mu\mu_1)$ . Then combining (37) and (38) together yields

$$\begin{aligned}
\|\mathbf{s}^{k-1} - \boldsymbol{\sigma}^*\| &= \|I - J_{k-1}Q_{k-1}\| \cdot \|J_{k-1}\mathbf{c}^{k-1} + \mathbf{w}^{k-1} - \boldsymbol{\sigma}^*\| \\
&\leq \|I - J_{k-1}Q_{k-1}\| \cdot \sqrt{n} \|U_{k-1}^T A(\mathbf{c}^{k-1})V_{k-1} - \Sigma^*\| \\
&\leq \sqrt{n}\mu^{2^k-2}\psi_1^{2^k} \leq \frac{\sqrt{n}}{\mu}(\mu\psi_1)^{2^k-1}\psi_1 \leq (\mu\mu_1\delta)^{2^k-1}\psi_1 \\
&< \mu\mu_1\delta \leq \min\{\frac{1}{2}, \frac{1}{2}\widehat{d}^*\}.
\end{aligned} \tag{39}$$

Thus one has

$$\frac{3}{2}\sigma_1^* > s_1^{k-1} > \frac{1}{2}(\sigma_1^* + \sigma_2^*) > s_2^{k-1} > \frac{1}{2}(\sigma_2^* + \sigma_3^*) > \cdots > \frac{1}{2}(\sigma_{n-1}^* + \sigma_n^*) > s_n^{k-1} > \frac{3}{4}\sigma_n^* > 0. \tag{40}$$

By (22) once again, we have  $\delta \leq d^*/(2\mu\mu_1(4\sigma_1^* + 1))$ . We further deduce from (39) that

$$\begin{aligned}
|(s_i^{k-1})^2 - (s_j^{k-1})^2| &\geq d^* - 4\sigma_1^* \|\mathbf{s}^{k-1} - \boldsymbol{\sigma}^*\| - 2\|\mathbf{s}^{k-1} - \boldsymbol{\sigma}^*\|^2 \\
&\geq d^* - (4\sigma_1^* + 1)\|\mathbf{s}^{k-1} - \boldsymbol{\sigma}^*\| \\
&\geq d^* - (4\sigma_1^* + 1)\mu\mu_1\delta \geq \frac{1}{2}d^*,
\end{aligned}$$

for any  $1 \leq i \neq j \leq n$ .

By (5) and Lemma 3.4, we obtain

$$\Phi_k = I + X_k + G_k, \quad \Psi_k = I + Y_k + H_k, \tag{41}$$

where  $\|G_k\| \leq \|X_k\|^2 \leq \psi_k^2$  and  $\|H_k\| \leq \|Y_k\|^2 \leq \psi_k^2$ . Using  $U_k = U_{k-1}\Phi_k$ ,  $V_k = V_{k-1}\Psi_k$ , and (41), we get by simple calculation,

$$U_k^T A(\mathbf{c}^k)V_k = S_{k-1} + R_k = \Sigma^* + (S_{k-1} - \Sigma^*) + R_k, \tag{42}$$

where

$$\begin{aligned}
R_k &= X_k(S_{k-1} - X_k S_{k-1} + S_{k-1} Y_k) Y_k - X_k^2 S_{k-1} - S_{k-1} Y_k^2 \\
&\quad + G_k^T (S_{k-1} + X_k S_{k-1} - S_{k-1} Y_k) (I + Y_k) \\
&\quad + (I - X_k + G_k^T) (S_{k-1} + X_k S_{k-1} - S_{k-1} Y_k) H_k.
\end{aligned}$$

Using (36), (40), (41), and the hypothesis on  $\psi_k$ , we have

$$\begin{aligned}
\|R_k\| &\leq 3\left(\frac{3}{2}\sigma_1^*\right)\|X_k\| \cdot \|Y_k\| + \frac{3}{2}\sigma_1^*\|X_k\|^2 + \frac{3}{2}\sigma_1^*\|Y_k\|^2 + 6\left(\frac{3}{2}\sigma_1^*\right)\|X_k\|^2 + 9\left(\frac{3}{2}\sigma_1^*\right)\|Y_k\|^2 \\
&\leq \frac{3}{2}\left(\frac{3}{2}\sigma_1^*\right)(\|X_k\|^2 + \|Y_k\|^2) + 10\left(\frac{3}{2}\sigma_1^*\right)(\|X_k\|^2 + \|Y_k\|^2) \\
&= \frac{23}{2}\left(\frac{3}{2}\sigma_1^*\right)\psi_k^2 \leq \mu_2\psi_k^2 \leq \mu_2\mu^{2^k-2}\psi_1^{2^k}.
\end{aligned} \tag{43}$$

Then by (39), (42), and (43),

$$\begin{aligned}
\|U_k^T A(\mathbf{c}^k)V_k - \Sigma^*\| &\leq \|S_{k-1} - \Sigma^*\| + \|R_k\| \leq \|\mathbf{s}^{k-1} - \boldsymbol{\sigma}^*\| + \|R_k\| \\
&\leq (\mu_2 + \sqrt{n})\mu^{2^k-2}\psi_1^{2^k} \leq \mu^{2^k-1}\psi_1^{2^k}.
\end{aligned} \tag{44}$$

We can conclude that (15) holds for all  $k \geq 1$ .

By (22), we have  $\delta \leq \delta_3$  and  $\delta \leq 3\delta_1/(16\mu\mu_1)$ . Then by Theorem 3.6, (24), and the hypothesis on  $V_k$ ,

$$\begin{aligned}\|V_k - V_*\| &\leq \sum_{j=2}^k \|V_j - V_{j-1}\| + \|V_1 - V_0\| + \|V_0 - V_*\| \\ &\leq \sum_{j=2}^k 2(\mu\psi_1)^{2^{j-1}} + 2\psi_1 + \frac{\delta_1}{4} \leq \frac{2(\mu\psi_1)^2}{1-(\mu\psi_1)^2} + \frac{\delta_1}{2} + \frac{\delta_1}{4} \\ &\leq \frac{4}{3}\mu\psi_1 + \frac{\delta_1}{2} + \frac{\delta_1}{4} \leq \frac{\delta_1}{4} + \frac{\delta_1}{2} + \frac{\delta_1}{4} = \delta_1.\end{aligned}$$

Similarly, we can prove that  $\|U_1^k - U_1^*\| \leq \delta_1$ . Thus  $\|J_k^{-1}\| \leq C$ .

As the proof of (16) for  $k = 1$ , we can obtain

$$\|I - J_k Q_k\| \leq 2\|I - J_{k-1} Q_{k-1}\|^2 + 2\|J_k - J_{k-1}\|^2 \|Q_{k-1}\|^2$$

and

$$\|J_k - J_{k-1}\| \leq n(\|U_k - U_{k-1}\| + \|V_k - V_{k-1}\|) \cdot \max_j \|A_j\|.$$

Thus by (37) and the hypothesis,

$$\|I - J_k Q_k\| \leq (2 + \mu_0)\mu^{2^k-2}\psi_1^{2^k} \leq \mu^{2^k-1}\psi_1^{2^k} \leq \psi_1 < 1 \quad (45)$$

and hence

$$\|Q_k\| \leq \|J_k^{-1}\| \cdot \|J_k Q_k\| \leq \|J_k^{-1}\|(1 + \|I - J_k Q_k\|) \leq 2C,$$

i.e., (16)–(17) hold for all  $k \geq 1$ .

Now, we have by Algorithm 1, (42), and (44),

$$\begin{aligned}\|\mathbf{c}^{k+1} - \mathbf{c}^k\| &\leq \|Q_k\| \cdot \|J_k \mathbf{c}^k + \mathbf{w}^k - \boldsymbol{\sigma}^*\| \leq 2C\sqrt{n}\|U_k^T A(\mathbf{c}^k)V_k - \Sigma^*\| \\ &\leq 2C\sqrt{n}(\mu_2 + \sqrt{n})\mu^{2^k-2}\psi_1^{2^k} \leq \mu_3\mu^{2^k-2}\psi_1^{2^k} \leq \mu^{2^k-1}\psi_1^{2^k}\end{aligned} \quad (46)$$

and then (18) holds for all  $k \geq 1$ .

By (22), one has  $\delta \leq 1/(2\mu\mu_1)$  and  $\delta \leq \widehat{d}^*/(2\mu\mu_1)$ . Then relations (23), (42), (44), and (45) imply that

$$\begin{aligned}\|\mathbf{s}^k - \boldsymbol{\sigma}^*\| &\leq \|I - J_k Q_k\| \cdot \|J_k \mathbf{c}^k + \mathbf{w}^k - \boldsymbol{\sigma}^*\| \leq \sqrt{n}\|U_k^T A(\mathbf{c}^k)V_k - \Sigma^*\| \\ &\leq \sqrt{n}(\mu_2 + \sqrt{n})\mu^{2^k-2}\psi_1^{2^k} \leq \mu_3\mu^{2^k-2}\psi_1^{2^k} \leq \mu^{2^k-1}\psi_1^{2^k} \\ &= (\mu\psi_1)^{2^k-1}\psi_1 \leq (\mu\mu_1\delta)^{2^k-1}\psi_1 < \mu\mu_1\delta \leq \min\{\frac{1}{2}, \frac{1}{2}\widehat{d}^*\}.\end{aligned} \quad (47)$$

Thus one has

$$\frac{3}{2}\sigma_1^* > s_1^k > \frac{1}{2}(\sigma_1^* + \sigma_2^*) > s_2^k > \frac{1}{2}(\sigma_2^* + \sigma_3^*) > \cdots > \frac{1}{2}(\sigma_{n-1}^* + \sigma_n^*) > s_n^k > \frac{3}{4}\sigma_n^* > 0.$$

By (22) again, one has  $\delta \leq d^*/(2\mu\mu_1(4\sigma_1^* + 1))$ . We further derive from (47) that

$$\begin{aligned}|(s_i^k)^2 - (s_j^k)^2| &\geq d^* - 4\sigma_1^*\|\mathbf{s}^k - \boldsymbol{\sigma}^*\| - 2\|\mathbf{s}^k - \boldsymbol{\sigma}^*\|^2 \\ &\geq d^* - (4\sigma_1^* + 1)\|\mathbf{s}^k - \boldsymbol{\sigma}^*\| \\ &\geq d^* - (4\sigma_1^* + 1)\mu\mu_1\|\mathbf{c}^0 - \mathbf{c}^*\| \geq \frac{1}{2}d^*,\end{aligned}$$

for any  $1 \leq i \neq j \leq n$ .

Relation (44) implies that

$$U_k^T A(\mathbf{c}^k) V_k - \Sigma^* = \widehat{R}_k, \quad \|\widehat{R}_k\| \leq (\mu_2 + \sqrt{n}) \mu^{2^k-2} \psi_1^{2^k} \leq \mu^{2^k-1} \psi_1^{2^k}. \quad (48)$$

Combining (48) with (6) gives rise to

$$X_{k+1} S_k - S_k Y_{k+1} = U_k^T (A(\mathbf{c}^{k+1}) - A(\mathbf{c}^k)) V_k + \widehat{R}_k - (S_k - \Sigma^*),$$

and we have by Lemma 3.5 and (46)–(48),

$$\begin{aligned} \psi_{k+1} = \sqrt{\|X_{k+1}\|^2 + \|Y_{k+1}\|^2} &\leq \sqrt{2\rho} (\|A(\mathbf{c}^{k+1}) - A(\mathbf{c}^k)\| + \|\widehat{R}_k\| + \|S_k - \Sigma^*\|) \\ &\leq \sqrt{2\rho} (\zeta \|\mathbf{c}^{k+1} - \mathbf{c}^k\| + \|\widehat{R}_k\| + \|\mathbf{s}^k - \boldsymbol{\sigma}^*\|) \\ &\leq \sqrt{2\rho} ((\zeta + 1)\mu_3 + \mu_2 + \sqrt{n}) \mu^{2^k-2} \psi_1^{2^k} \\ &\leq \mu_4 \mu^{2^k-2} \psi_1^{2^k} \leq \mu^{2^k-1} \psi_1^{2^k}. \end{aligned} \quad (49)$$

This verifies that (19) holds for all  $k \geq 1$ .

By Lemma 3.4 and (49), we obtain

$$\|U_{k+1} - U_k\| \leq 2\|X_{k+1}\| \leq 2\psi_{k+1} \leq 2\mu^{2^k-1} \psi_1^{2^k}$$

and

$$\|V_{k+1} - V_k\| \leq 2\|Y_{k+1}\| \leq 2\psi_{k+1} \leq 2\mu^{2^k-1} \psi_1^{2^k}.$$

Therefore, the estimates (20) and (21) hold for all  $k \geq 1$ .  $\square$

Finally, we show that our method converges at least quadratically in the root sense. We first recall the definition of convergence rate in the root sense [16, Chapter 9].

**Definition 3.8** Let  $\{\mathbf{x}^k\}$  be a sequence to a limit  $\mathbf{x}^*$ . Then the numbers

$$R_p\{\mathbf{x}^k\} = \begin{cases} \limsup_{k \rightarrow \infty} \|\mathbf{x}^k - \mathbf{x}^*\|^{1/k}, & \text{if } p = 1, \\ \limsup_{k \rightarrow \infty} \|\mathbf{x}^k - \mathbf{x}^*\|^{1/p^k}, & \text{if } p > 1, \end{cases}$$

are the convergence factors in the root sense of  $\{\mathbf{x}^k\}$ . The quantity

$$O_R(\mathbf{x}^*) = \begin{cases} \infty, & \text{if } R_p\{\mathbf{x}^k\} = 0, \quad \forall p \in [1, \infty), \\ \inf\{p \in [1, \infty) \mid R_p\{\mathbf{x}^k\} = 1\}, & \text{otherwise,} \end{cases}$$

is called **the convergence rate in the root sense of  $\{\mathbf{x}^k\}$  at  $\mathbf{x}^*$** .

The following theorem gives the main convergence result whose proof is similar to that of Theorems 2 and 3 in [4] (see also [5, Theorems 4.10 and 4.12]) and therefore we omit it.

**Theorem 3.9** Suppose that  $\sigma_1^* > \sigma_2^* > \dots > \sigma_n^* > 0$  and Jacobian matrix  $J(\mathbf{c}^*)$  is invertible. Then there exists a constant  $\epsilon > 0$  such that, when  $\|\mathbf{c}^0 - \mathbf{c}^*\| \leq \epsilon$ , the sequences  $\{\mathbf{c}^k\}$ ,  $\{X_k\}$ ,  $\{Y_k\}$ ,  $\{U_k\}$ ,  $\{V_k\}$ , and  $\{U_k^T A(\mathbf{c}^k) V_k\}$  generated by Algorithm 1 converge at least quadratically in the root sense.

## 4 Numerical Tests

In this section, we report some numerical tests to illustrate the effectiveness of our proposed method. We compare the numerical performance of Algorithm 1 with that of the inexact Newton method in [5]. All the tests are implemented in `Matlab` 7.0 on a PC Inter Pentium R of 3.00 GHz CPU.

For the completeness of our presentation, we recall the inexact Newton method [5] as follows.

### Algorithm 2: The Inexact Newton Method

- I. Given  $\gamma \in (1, 2]$  and  $\mathbf{c}^0 \in \mathbb{R}^n$ , compute the singular values  $\{\sigma_i^0 = \sigma_i(A(\mathbf{c}^0))\}_{i=1}^n$ , the normalized left singular vectors  $\{\mathbf{u}_j^0 = \mathbf{u}_j(A(\mathbf{c}^0))\}_{j=1}^m$ , and the normalized right singular vectors  $\{\mathbf{v}_i^0 = \mathbf{v}_i(A(\mathbf{c}^0))\}_{i=1}^n$  of  $A(\mathbf{c}^0)$ . Form Jacobian matrix  $J_0$  and  $\mathbf{w}^0 \in \mathbb{R}^n$  by (4) and apply an iterative method (e.g., the QMR method [17]) to solving Jacobian equation

$$J_0 \mathbf{c}^1 = \boldsymbol{\sigma}^* - \mathbf{w}^0$$

such that

$$\|J_0 \mathbf{c}^1 - \boldsymbol{\sigma}^* + \mathbf{w}^0\| \leq r^0 \|\boldsymbol{\sigma}^* - \mathbf{w}^0\|,$$

where  $r^0 = \|\boldsymbol{\sigma}^0 - \boldsymbol{\sigma}^*\|^\gamma / \|\boldsymbol{\sigma}^*\|^\gamma$  with  $\boldsymbol{\sigma}^0 = (\sigma_1^0, \dots, \sigma_n^0)^T$ .

- II. For  $k = 1, 2, \dots$ , until convergence, do:

(i)–(v) as II (i)–(v) in Algorithm 1.

(vi) Apply an iterative method to solving the approximate Jacobian equation

$$J_k \mathbf{c}^{k+1} = \boldsymbol{\sigma}^* - \mathbf{w}^k$$

such that

$$\|J_k \mathbf{c}^{k+1} - \boldsymbol{\sigma}^* + \mathbf{w}^k\| \leq r^k \|\boldsymbol{\sigma}^* - \mathbf{w}^k\|,$$

where

$$r^k = \|\boldsymbol{\sigma}^k - \boldsymbol{\sigma}^*\|^\gamma / \|\boldsymbol{\sigma}^*\|^\gamma, \quad \boldsymbol{\sigma}^k = (\sigma_1^k, \dots, \sigma_n^k)^T, \quad \sigma_i^k = (\mathbf{u}_i^k)^T A(\mathbf{c}^k) \mathbf{v}_i^k,$$

for  $1 \leq i \leq n$ .

For simplicity, in our numerical experiments, we focus on the following four cases: a)  $m = 100$  and  $n = 60$ ; b)  $m = 300$  and  $n = 120$ ; c)  $m = 600$  and  $n = 300$ ; d)  $m = 800$  and  $n = 400$ . For the ISVP, we first generate the basis matrices  $A_0, A_1, \dots, A_n$  and a solution  $\mathbf{c}^*$  randomly. Then we use the singular values of  $A(\mathbf{c}^*)$  as the given singular values  $\{\sigma_j^*\}_{j=1}^n$ . The initial guess  $\mathbf{c}^0$  is generated via perturbing each entry of  $\mathbf{c}^*$  uniformly distributed in an interval of  $[-\max_j |c_j^*| \cdot \beta, \max_j |c_j^*| \cdot \beta]$  for different  $\beta > 0$ .

For demonstration purpose, all linear systems appeared in Algorithms 1 and 2 were solved by the QMR method [17] via the `Matlab`-provided `QMR` function, where the maximal number of iteration is set to be 1000. In particular, for approximate Jacobian equations in Algorithm 2,

we use the preconditioned QMR method with the given stopping tolerance and adopt `Matlab`-provided Incomplete LU factorization as the preconditioner, i.e., `LUINC(A,drop-tolerance)`, where the drop tolerance is set to be 0.01. Also, the initial guess for the approximate Jacobian equation in the  $(k + 1)$ th outer iteration is set to be  $\mathbf{c}^k$  obtained at the  $k$ th outer iteration. The stopping tolerance for the other linear systems appeared in Algorithms 1 and 2 were set to be  $10^{-14}$  so that the desired solutions are obtained. Algorithms 1 and 2 stop when

$$\|U_k^T A(\mathbf{c}^k)V_k - \Sigma^*\|_F \leq 10^{-8}$$

and the maximal number of outer iterations is set to be 20.

Next, we report our numerical results. Tables 1–4 list the residual values for cases a)–d) with different choices of  $\beta$ . Here `it.` and  $\text{cond}_2(J_k)$  denote the number of the  $k$ th outer iteration ( $k = 0, 1, 2, \dots$ ) and the condition number of  $J_k$  at the  $k$ th outer iteration (For Algorithm 2,  $\text{cond}_2(J_k)$  means the condition number of  $J_k$  for  $\gamma = 2$ ), respectively. For simplicity, in Tables 3 and 4, we only report the numerical results for Algorithm 2 with  $\gamma = 2.0$ .

From Tables 1–4, we observe that both Algorithms 1 and 2 converge superlinearly in the root sense if approximate Jacobian equations are well-conditioned. However, when the condition numbers of approximate Jacobian matrices  $J_k$  become large, Algorithm 1 works much better than Algorithm 2.

Table 1: Residual values for case a)

| $\beta$ | it. | Algorithm 2    |                |                |              |                      | Algorithm 1  |                      |
|---------|-----|----------------|----------------|----------------|--------------|----------------------|--------------|----------------------|
|         |     | $\gamma = 1.5$ | $\gamma = 1.6$ | $\gamma = 1.8$ | $\gamma = 2$ | $\text{cond}_2(J_k)$ |              | $\text{cond}_2(J_k)$ |
| 0.001   | 0   | $8.08e - 1$    | $8.08e - 1$    | $8.08e - 1$    | $8.08e - 1$  | $3.95e + 2$          | $8.08e - 1$  | $3.95e + 2$          |
|         | 1   | $2.00e - 2$    | $1.83e - 2$    | $1.83e - 2$    | $1.83e - 2$  | $3.48e + 2$          | $1.83e - 2$  | $3.48e + 2$          |
|         | 2   | $6.89e - 5$    | $5.88e - 5$    | $5.88e - 5$    | $5.85e - 5$  | $3.50e + 2$          | $8.09e - 5$  | $3.50e + 2$          |
|         | 3   | $1.62e - 9$    | $1.05e - 9$    | $1.05e - 9$    | $1.03e - 9$  |                      | $6.18e - 8$  | $3.50e + 2$          |
|         | 4   |                |                |                |              |                      | $2.94e - 12$ |                      |
| 0.0001  | 0   | $8.10e - 2$    | $8.10e - 2$    | $8.10e - 2$    | $8.10e - 2$  | $3.54e + 2$          | $8.10e - 2$  | $3.54e + 2$          |
|         | 1   | $1.77e - 4$    | $1.77e - 4$    | $1.77e - 4$    | $1.77e - 4$  | $3.50e + 2$          | $1.77e - 4$  | $3.50e + 2$          |
|         | 2   | $5.71e - 9$    | $5.59e - 9$    | $5.59e - 9$    | $5.04e - 9$  |                      | $6.43e - 9$  |                      |

## 5 Concluding Remarks

In this paper, we propose a Ulm-like method for solving inverse singular value problems. This method avoids solving approximate Jacobian equations in Newton’s methods. Under some mild assumptions, we show that our method converges at least quadratically in the root sense. Numerical results illustrate the effectiveness of our method. In our proof, however, we assume that all the given singular values are positive and distinct. An interesting topic is to extend the proposed method to the cases of multiple singular values and of zero singular values, which needs further investigation.

Table 2: Residual values for case b)

| $\beta$  | it.       | Algorithm 2    |                |                |              |                      | Algorithm 1 |                      |
|----------|-----------|----------------|----------------|----------------|--------------|----------------------|-------------|----------------------|
|          |           | $\gamma = 1.5$ | $\gamma = 1.6$ | $\gamma = 1.8$ | $\gamma = 2$ | $\text{cond}_2(J_k)$ |             | $\text{cond}_2(J_k)$ |
| 0.001    | 0         | $3.96e+0$      | $3.96e+0$      | $3.96e+0$      | $3.96e+0$    | $9.84e+4$            | $3.96e+0$   | $9.84e+4$            |
|          | 1         | $1.00e-1$      | $6.97e-2$      | $5.13e-2$      | $5.13e-2$    | $7.24e+4$            | $3.25e-2$   | $1.38e+5$            |
|          | 2         | $1.28e-2$      | $2.08e-2$      | $1.44e-2$      | $1.44e-2$    | $1.24e+5$            | $8.10e-3$   | $1.17e+5$            |
|          | 3         | $4.33e-3$      | $8.56e-3$      | $1.22e-2$      | $1.22e-2$    | $1.27e+5$            | $2.04e-5$   | $1.18e+5$            |
|          | 4         | $1.76e-3$      | $7.73e-3$      | $6.86e-3$      | $6.86e-3$    | $1.27e+5$            | $1.58e-8$   | $1.18e+5$            |
|          | 5         | $1.20e-3$      | $5.27e-3$      | $6.86e-3$      | $6.86e-3$    | $1.27e+5$            | $4.83e-12$  |                      |
|          | $\vdots$  | $\vdots$       | $\vdots$       | $\vdots$       | $\vdots$     | $\vdots$             |             |                      |
| 20       | $5.36e-4$ | $9.92e-4$      | $6.86e-3$      | $6.86e-3$      | $1.27e+5$    |                      |             |                      |
| 0.0001   | 0         | $3.96e-1$      | $3.96e-1$      | $3.96e-1$      | $3.96e-1$    | $1.16e+5$            | $3.96e-1$   | $1.16e+5$            |
|          | 1         | $1.95e-2$      | $1.95e-2$      | $1.95e-2$      | $1.95e-2$    | $1.23e+5$            | $2.48e-4$   | $1.18e+5$            |
|          | 2         | $5.09e-3$      | $5.09e-3$      | $5.09e-3$      | $5.09e-3$    | $1.10e+5$            | $1.06e-6$   | $1.18e+5$            |
|          | 3         | $1.19e-3$      | $1.19e-3$      | $1.19e-3$      | $1.19e-3$    | $1.11e+5$            | $4.98e-11$  |                      |
|          | $\vdots$  | $\vdots$       | $\vdots$       | $\vdots$       | $\vdots$     | $\vdots$             |             |                      |
|          | 20        | $8.39e-4$      | $8.39e-4$      | $8.39e-4$      | $8.39e-4$    | $1.12e+5$            |             |                      |
|          | 0.00001   | 0              | $3.96e-2$      | $3.96e-2$      | $3.96e-2$    | $3.96e-2$            | $1.17e+5$   | $3.96e-2$            |
| 1        | $8.44e-4$ | $8.44e-4$      | $8.44e-4$      | $8.44e-4$      | $1.17e+5$    | $2.46e-6$            | $1.18e+5$   |                      |
| 2        | $4.79e-4$ | $4.79e-4$      | $4.79e-4$      | $4.79e-4$      | $1.18e+5$    | $1.10e-10$           |             |                      |
| $\vdots$ | $\vdots$  | $\vdots$       | $\vdots$       | $\vdots$       | $\vdots$     |                      |             |                      |
| 20       | $4.77e-6$ | $4.77e-6$      | $4.77e-6$      | $4.77e-6$      | $1.18e+5$    |                      |             |                      |

**Acknowledgments.** We are very grateful to the editor and two anonymous referees for their valuable comments which have considerably improved this paper.

## References

- [1] Z. J. Bai, *Inexact Newton methods for inverse eigenvalue problems*, Appl. Math. Comput., 172 (2006), pp. 682–689.
- [2] Z. J. Bai, R. H. Chan, and B. Morini, *An inexact Cayley transform method for inverse eigenvalue problems*, Inverse Problems, 20 (2004), pp. 1675–1689.
- [3] Z. J. Bai and X. Q. Jin, *A note on the Ulm-like method for inverse eigenvalue problems*, Recent Advances in Scientific Computing and Matrix Analysis, pp. 1–7. Eds: X. Q. Jin, H. W. Sun, and S. W. Vong, International Press, Somerville; and Higher Education Press, Beijing, 2011.
- [4] Z. J. Bai, B. Morini, and S. F. Xu, *On the local convergence of an iterative approach for inverse singular value problems*, J. Comput. Appl. Math., 198 (2007), pp. 344–360.



Table 3: Residual values for case c)

| $\beta$ | it.      | Algorithm 2  |                      | Algorithm 1 |                      |
|---------|----------|--------------|----------------------|-------------|----------------------|
|         |          | $\gamma = 2$ | $\text{cond}_2(J_k)$ |             | $\text{cond}_2(J_k)$ |
| 0.0001  | 0        | $3.30e-1$    | $2.31e+5$            | $3.30e-1$   | $2.31e+5$            |
|         | 1        | $1.00e-1$    | $2.48e+5$            | $2.81e-3$   | $2.21e+5$            |
|         | 2        | $1.00e-1$    | $2.15e+5$            | $4.28e-6$   | $2.20e+5$            |
|         | 3        | $1.00e-1$    | $1.92e+5$            | $9.77e-11$  |                      |
|         | $\vdots$ | $\vdots$     | $\vdots$             |             |                      |
|         | 20       | $1.00e-1$    | $1.84e+5$            |             |                      |
| 0.00001 | 0        | $3.30e-2$    | $2.21e+5$            | $3.30e-2$   | $2.21e+5$            |
|         | 1        | $2.68e-2$    | $2.23e+5$            | $2.89e-5$   | $2.20e+5$            |
|         | 2        | $2.48e-2$    | $2.20e+5$            | $4.63e-10$  |                      |
|         | $\vdots$ | $\vdots$     | $\vdots$             |             |                      |
|         | 20       | $1.45e-2$    | $2.16e+5$            |             |                      |

- [5] Z. J. Bai and S. F. Xu, *An inexact Newton-type method for inverse singular value problems*, Linear Algebra Appl., 429 (2008), pp. 527–547.
- [6] R. H. Chan, H. L. Chung, and S. F. Xu, *The inexact Newton-like method for inverse eigenvalue problem*, BIT, 43 (2003), pp. 7–20.
- [7] D. L. Chu and M. T. Chu, *Low rank update of singular values*, Math. Comp., 75 (2006), pp. 1351–1366.
- [8] M. T. Chu, *Numerical methods for inverse singular value problems*, SIAM J. Numer. Anal., 29 (1992), pp. 885–903.
- [9] M. T. Chu, *Inverse eigenvalue problems*, SIAM Rev., 40 (1998), pp. 1–39.
- [10] M. T. Chu and G. H. Golub, *Structured inverse eigenvalue problems*, Acta Numer., 11 (2002), pp. 1–71.
- [11] M. T. Chu and G. H. Golub, *Inverse Eigenvalue Problems: Theory, Algorithms, and Applications*, Oxford University Press, Oxford, 2005.
- [12] S. Friedland, J. Nocedal, and M. L. Overton, *The formulation and analysis of numerical methods for inverse eigenvalue problems*, SIAM J. Numer. Anal., 24 (1987), pp. 634–667.
- [13] G. H. Golub and C. F. Van Loan, *Matrix Computations*, 3rd edition, Johns Hopkins University Press, Baltimore, 1996.
- [14] E. Montaña, M. Salas, and R. L. Soto, *Nonnegative matrices with prescribed extremal singular values*, Comput. Math. Appl., 56 (2008), pp. 30–42.

Table 4: Residual values for case d)

| $\beta$  | it.      | Algorithm 2  |                      | Algorithm 1  |                      |
|----------|----------|--------------|----------------------|--------------|----------------------|
|          |          | $\gamma = 2$ | $\text{cond}_2(J_k)$ |              | $\text{cond}_2(J_k)$ |
| 0.00001  | 0        | $1.18e - 1$  | $4.63e + 6$          | $1.18e - 1$  | $4.63e + 6$          |
|          | 1        | $8.18e - 2$  | $1.57e + 7$          | $6.68e - 5$  | $3.25e + 6$          |
|          | 2        | $7.68e - 2$  | $1.16e + 6$          | $6.58e - 6$  | $3.84e + 6$          |
|          | 3        | $7.27e - 2$  | $3.16e + 6$          | $5.72e - 7$  | $4.04e + 6$          |
|          | 4        | $6.35e - 2$  | $2.35e + 6$          | $7.25e - 9$  |                      |
|          | $\vdots$ | $\vdots$     | $\vdots$             |              |                      |
|          | 20       | $4.87e - 2$  | $3.70e + 6$          |              |                      |
| 0.000001 | 0        | $1.18e - 2$  | $4.11e + 6$          | $1.18e - 2$  | $4.11e + 6$          |
|          | 1        | $8.15e - 3$  | $6.00e + 6$          | $5.57e - 7$  | $4.05e + 6$          |
|          | 2        | $7.44e - 3$  | $3.60e + 6$          | $9.06e - 10$ |                      |
|          | $\vdots$ | $\vdots$     | $\vdots$             |              |                      |
|          | 20       | $6.26e - 3$  | $3.55e + 6$          |              |                      |

- [15] E. Montañó, M. Salas, and R. L. Soto, *Positive matrices with prescribed singular values*, *Proyecciones*, 27 (2008), pp. 289–305.
- [16] J. M. Ortega and W. C. Rheinboldt, *Iterative Solution of Nonlinear Equations in Several Variables*, Academic Press, 1970.
- [17] Y. Saad, *Iterative Methods for Sparse Linear Systems*, PWS Publishing Company, Boston, 1996.
- [18] W. P. Shen and C. Li, *A Ulm-like Cayley transform method for inverse eigenvalue problems*, Preprint, Department of Mathematics, Zhejiang University, P. R. China, 2010.
- [19] W. P. Shen, C. Li, and X. Q. Jin, *A Ulm-like method for inverse eigenvalue problems*, *Appl. Numer. Math.*, 61 (2011), pp. 356–367.
- [20] J. A. Tropp, I. S. Dhillon, and R. W. Heath Jr., *Finite-step algorithms for constructing optimal CDMA signature sequences*, *IEEE Trans. Inform. Theory*, 50 (2004), pp. 2916–2921.
- [21] S. Ulm, *On iterative methods with successive approximation of the inverse operator*, *Izv. Akad. Nauk Est. SSR* 16 (1967), pp. 403–411. (In Russian)
- [22] S. F. Xu, *An Introduction to Inverse Eigenvalue Problems*, Peking University Press and Vieweg Publishing, Beijing, 1998.